

ORIGINAL RESEARCH

Predicting operative difficulties during laparoscopic cholecystectomy using clinical and imaging criteria in pre-operative period

¹Dr. Abhijit D Hiregoudar, ²Dr. Sandeep Kumar R

¹Associate Professor, Department of General Surgery, KMCRI, Hubli, Karnataka, India

²Post Graduate, Department of General Surgery, KMCRI, Hubli, Karnataka, India

Corresponding Author

Dr. Abhijit D Hiregoudar

Associate Professor, Department of General Surgery, KMCRI, Hubli, Karnataka, India

Received: 19Jan, 2025

Accepted: 20Feb, 2025

ABSTRACT

Introduction: Laparoscopic cholecystectomy has become the procedure of choice for the management of symptomatic gallstone disease. Difficulty in laparoscopic cholecystectomy is often encountered by surgeons and is associated with complications and a higher conversion rate. This study aims at identifying difficult laparoscopic cholecystectomies by clinical and imaging assessment and determining the utility of a pre-operative difficulty scoring system based on intra-operative findings. **Patients and Methods:** A hospital based observational study was conducted including all patients above age 18 years undergoing laparoscopic cholecystectomy. Randhawa's preoperative score was calculated for every patient based on history, clinical and imaging parameters and the degree of operative difficulty was predicted. Another scoring was done using Sugrue's scoring system based on intra-operative findings. The outcome factors studied were degree of operative difficulty or need for conversion and intra-operative and post-operative complications. The statistical analysis was done by using Statistical software R version 4.4.0 and Microsoft excel. **Results:** A total 98 patients with mean age 47.39 ±14.34 years with 71 females and 27 males were enrolled. Of them 47 (47.96%) had difficult laparoscopic cholecystectomies. The conversion rate to open cholecystectomy was 8.5%. Age above 50 years, prior history of acute cholecystitis, thickened gall bladder wall and stone impaction of Hartmann's pouch were statistically significant pre-operative risk factors. The sensitivity of the preoperative score in identifying difficult cases was 88.24% with the specificity of 95.74%; positive predictive value of 95.74% and the negative predictive value of 88.24% and the accuracy was 92.86%. **Conclusion:** Pre-operative score and intra-operative findings are helpful in identifying difficult laparoscopic cholecystectomy.

Key words: Difficult laparoscopic cholecystectomy, predictive factors for difficult cholecystectomy

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

1. INTRODUCTION

Laparoscopic cholecystectomy has become the procedure of choice for the management of symptomatic gallstone disease. Advantages of laparoscopic cholecystectomy compared with open cholecystectomy are well described and usually include decreased postoperative pain, shorter ileus, earlier oral intake, and earlier return to normal activities with better cosmesis¹. "Difficult gallbladder" (DGB) is a term coined to denote a procedure with an increased surgical risk compared to standard cholecystectomies and has been associated with difficult dissection, altered anatomy and increased risk of bleeding^{2,3}. It is reported with an incidence of up to 26% in large series and a conversion rate of up to

26.5%⁴. There are many risk factors which make laparoscopic surgery difficult like old age, male gender, attacks of acute cholecystitis with fever and leukocytosis, obesity, previous abdominal surgery, clinical signs of acute cholecystitis, and some specific ultrasonographic findings i.e. thickened gall bladder wall, distended gall bladder, pericholecystic fluid collection, and impacted stone⁵.

Preoperative complexity estimation helps surgeons decide whether to proceed with a minimally invasive approach or perform an open procedure⁶. The use of a predictive score of operative difficulty is thus of primary interest to identify high-risk procedures and could be helpful to improve patient counselling, optimize surgical planning and operating room

efficiency, detect patients at risk of complications and change, when necessary, the operative technique. This study aims at identifying difficult laparoscopic cholecystectomies by clinico-imaging assessment and also at assessing the utility of Randhawa's pre-operative scoring based on intra-operative findings and final outcome of conversion rate and complications.

2. PATIENTS AND METHODS

An observational study was carried out at a tertiary care hospital after obtaining ethical clearance from

institutional ethics committee. All patients undergoing laparoscopic cholecystectomy for various indications such as cholelithiasis, acalculous cholecystitis, gall bladder polyps or porcelain gall bladder, were included in this study. Patients <18 years in age, unfit for anesthesia and those not consenting to join the study were excluded. A preoperative scoring was done for every patient; based on the demographic clinical and imaging findings and the degree of operative difficulty was predicted using the scoring system introduced by Randhawa *et al.*⁷[Table 1].

Table 1: Preoperative (Randhawa) scoring system⁷

Scoring Factors		Score	Maximum Score
History	Age	<50 years	0
		> 50 years	1
	Gender	Male	1
		Female	0
	History of hospitalization for acute cholecystitis	Yes	4
		NO	0
Clinical parameters	BMI	<25	0
		25-27.5	1
	Abdominal scar	>27.5	2
		No	0
		Infraumbilical	1
		Supraumbilical	2
	Palpable gall bladder	Yes	1
		No	0
Radiological parameters	Gall bladder wall thickness	Thin <4mm	0
		Thick ≥ 4mm	2
	Pericholecystic collection	No	0
		Yes	1
	Impacted stone in Hartmann's pouch	NO	0
		Yes	1
Total maximum score			15

Score up to 5 predicted easy, 6-10 difficult and >10 a very difficult laparoscopic cholecystectomy. The degree of operative difficulty was assessed based

on the laparoscopic intra-operative findings using the scoring system introduced by Sugrue *et al.*⁸[TABLE 2].

Table 2: Intra-operative Grading system for Difficult Laparoscopic Cholecystectomy (Sugrue *et al.*)⁸

Scoring factors	Score	Maximum score
Gall bladder appearance	Adhesions < 50% of gall bladder	1
	Adhesions burying the gall bladder	3
Distension/Contraction	Distended or Contracted shriveled Gall Bladder	1
	Unable to grasp the gall bladder with atraumatic laparoscopic forceps	1
Access	Stone of 1 cm or > 1 cm impacted in the Hartman's pouch	1
	BMI > 30	1
Severe Sepsis/Complication	Adhesions from previous surgery limiting access	1
	Bile or Pus outside gall bladder	1
	Time to identify cystic artery and cystic duct > 90 minutes	1
Total score		10

In this scoring system a score of <2 was considered easy, 2 to 4 moderately difficult, 5-7 very difficult, and 8 to 10, extremely difficult. The scores were compared in each patient to assess the reliability of pre-operative score. The outcome factors studied were degree of operative difficulty or need for conversion and intra-operative and post-operative complications. The statistical analysis was done by Statistical software R version 4.4.0. Univariate and multivariate regression analysis was carried out. Kaplan Maier survivor analysis was also done to find out the cut off levels of pre-operative and intra-operative scoring.

3. OBSERVATIONS AND RESULTS

A total of 98 patients who underwent laparoscopic cholecystectomy were enrolled in present study. Their mean age was 47.39 ± 14.34 years with a range of 18 to 80 years. There was female preponderance with 71 females as compared to 28 males and a male: female ratio of 1:2.62. Difficult cholecystectomy was labelled in 47(47.96%) cases based on intra-operative criteria mentioned in methodology. Of these 47 patients, 4(8.5%) required conversion to open cholecystectomy.

With an aim to identify predictors of difficult laparoscopic cholecystectomy a sub group analysis was carried out by dividing the patients into difficult cholecystectomy group and easy cholecystectomy group based on the intra-operative findings. Age was found to be one of the risk factors with 31 (65.96%) out of the 47 patients above the age of 50 years undergoing a difficult laparoscopic cholecystectomy and the mean age of cases labelled as difficult was 50.29 ± 12.298 years as compared to 39.3 ± 13.974 years in the cases labelled as easy. This was statistically significant with p value < 0.01. On evaluating gender as a risk factor, it was observed that 12 (44.44%) out of 27 males underwent a difficult laparoscopic cholecystectomy as compared to 35(49.3%) out of the 71 females. Although numerically it appears that male gender is a risk factor for difficult laparoscopic cholecystectomy, however statistical significance was not found [p value >0.05]. On assessing the role of BMI as a risk factor it was noted that 39 (60%) out of the 65 patients that had a BMI greater than 25, underwent a difficult laparoscopic cholecystectomy. Statistical significance of BMI as a risk factor was found with [p value <0.01]. The history of hospitalization for prior attack of acute cholecystitis was inquired for and noted in 12(12.25%) patients. Of them, 11 (91.67%) underwent a difficult laparoscopic cholecystectomy. This factor was found to be statistically significant with p value < 0.01. History of undergoing abdominal surgery in past was inquired and presence of an abdominal scar was looked for and noted during examination and was found to be present in 27 patients; of which 12(44.45%) underwent difficult laparoscopic cholecystectomy. However, history of previous

abdominal surgery was not found to be a predictor of difficult laparoscopic cholecystectomy [p value > 0.05]. None of the 98 subjects that underwent laparoscopic cholecystectomy had a palpable gall bladder which was supposed to be one of the predictors of difficult laparoscopic cholecystectomy. On evaluating the patients using imaging modalities it was observed that thickened gall bladder which was suggestive of chronic cholecystitis was seen in 52(53.06%) patients. Of them 43 (82.69%) underwent a difficult laparoscopic cholecystectomy. Hence Thickened gall bladder wall was a statistically significant predictor of difficult laparoscopic cholecystectomy with p value < 0.01. Another imaging finding that is pericholecystic collection was seen in 24 (24.49%) patients. Of them 22(91.66%) underwent a difficult laparoscopic cholecystectomy. It was a statistically significant predictor [p value <0.01]. Stone impaction of Hartmann's pouch was noticed in 17(17.35%) cases out of which 16(94.12%) underwent a difficult laparoscopic cholecystectomy. This was statistically significant with p value < 0.01. On evaluating the intra-operative findings based on which the difficulty of laparoscopic cholecystectomy was graded it was observed that; gall bladder adhesions, distended or contracted gall bladder, inability to grasp the gall bladder using atraumatic laparoscopic forceps, stone impaction of Hartmann's pouch, time taken to identify and ligate the cystic artery > 90 minutes and intra-operative spillage of gall bladder contents were harbingers of difficult laparoscopic cholecystectomy with a p value < 0.01.[Table 3]. Intra-abdominal adhesions from previous surgeries and obesity which can limiting the access to gall bladder were not found to be associated with difficult laparoscopic cholecystectomy with statistical significance.

In cases of conventional laparoscopic cholecystectomy, the mean operative time of those which were graded as easy was 64.02 ± 10.38 minutes and range was between 40 to 95 minutes whereas in cases found to be difficult the operative time ranged between 75 to 150 minutes with a mean of 110 ± 19.003 minutes. This was statistically significant with p value < 0.01 suggesting difficult procedure obviously needed longer duration for completion of procedure. On evaluating the Randhawa's pre-operative score based on the intra-operative findings, it was found that easy cholecystectomies scored 2.52 ± 1.81 as compared to 7.25 ± 2.22 in the difficult group. The sensitivity of the preoperative score in predicting difficult cases was 85.7% with the specificity of 96.43%; positive predictive value of 92.31% and the negative predictive value of 93.10%. The accuracy of the pre-operative scoring system was 92.86%. It was observed that the subjects that were labelled to have undergone easy laparoscopic cholecystectomy the mean intra-operative score (Sugrue *et al.*) was 1.32 ± 0.956 and all the subjects that were labelled to have undergone

difficult laparoscopic cholecystectomy the mean score was 5.79 ± 1.067 a multivariate analysis was carried out of the factors found to be significant by univariate

analysis and it was observed that no singular factor was independently identifiable with difficult laparoscopic cholecystectomy.

Table 3: Showing univariate analysis of intra-operative parameters

Serial Number	Patient characteristics (N=84)	Intra-operative outcome based on grading		Univariate analysis P value
		Difficult	Easy	
1.	Amount of gall bladder adhesions	> 50%	13(100%)	<0.01
		< 50%	14(50%)	
		No adhesions	42(97.67%)	
2.	Distended/Contracted gallbladder	Yes	20(47.62%)	<0.005
		No	8(19.05%)	
3.	Inability to grasp the gall bladder with traumatic laparoscopic forceps	Yes	22(68.75%)	<0.01
		No	6(11.54%)	
4.	Intra-operative spillage of gallbladder contents	Yes	22(81.48%)	<0.01
		No	6(10.53%)	
5.	Time taken to identify cystic artery	> 90 minutes	19(86.36%)	<0.01
		< 90 minutes	9(13.85%)	

On carrying out Kaplan-Meier survival analysis and plotting a receiver operating curve of pre-operative score and intra-operative score; the pre-operative cut off score for difficult laparoscopic cholecystectomy was found to be 6 or more [Figure 1] and the intra-operative cut off score was 5 or more, [Figure 2]

suggesting that any pre-operative score of 6 or more was associated with difficult laparoscopic cholecystectomy while intra-operative score of 5 or more was suggestive of difficult laparoscopic cholecystectomy.




Figure 1: ROC of pre-operative score




Figure 2: ROC of intra-operative score

4. DISCUSSION

Cholecystectomy is currently one of the commonest reasons for admission to hospital with an associated mortality of 0.45 to 6% depending on severity of gallbladder disease⁹. With technological advancement and increase in the expertise, the complication rate has reached a remarkably low level at 2.0-6.0%¹⁰. Current literature however, suggests that the rate of intra-operative conversion from laparoscopic cholecystectomy to open cholecystectomy is 1%-15%¹¹⁻¹³. Difficult laparoscopic cholecystectomy still, is associated with serious operative and postoperative complications and a relatively high conversion rate¹⁴. Various “damage control” procedures, such as cholecystostomy (percutaneous or trocar site), fundus first approach, subtotal cholecystectomy and more recently, an “inside approach” have been proposed to reduce the risk of intra-operative complication in cases of difficult laparoscopic cholecystectomy¹⁵.

Age above 50 years proved to be a significant risk factor for difficult laparoscopic cholecystectomy in this study. It is consistent with results in literature¹⁶.

The reason for older age being at risk of difficult laparoscopic cholecystectomy is due to a longer history of gallstones and increased number of acute attacks of cholecystitis. Besides, elderly patients have a higher likelihood of complicated biliary pathology¹⁷⁻¹⁹. Males are at a higher risk of difficult gall bladder¹⁻

²⁰. Men with symptomatic gall bladder are more prone to inflammation and fibrosis with the same disease intensity thus leading to difficulty in dissection. They obtained higher levels of collagen, hydroxyproline, macrophages, mast cells and eosinophils in gallbladder wall and pericholecystic tissue which might explain the male fibrogenic propensity²¹. Also, men tend to present late as they pay less attention to subtle symptoms, therefore, may have more advanced disease²². However, in present study no statistical significance was found to co-relate male gender with difficult laparoscopic cholecystectomy and these results are quite consistent with literature^{8, 23}. BMI >30 to be significantly associated with difficulty in umbilical port entry and creating pneumoperitoneum^{24, 25}. Present study did not identify raised BMI as a risk factor for difficult laparoscopic cholecystectomy. These results are consistent with some studies²³. However, other studies did recognize raised BMI as a risk factor²⁶.

In present study history of prior attacks of acute cholecystitis was identified as a risk factor for difficult laparoscopic cholecystectomy which is quite consistent with literature²⁷⁻²⁹. Patient, who require hospitalization for repeated attacks of acute cholecystitis, carry more chances of difficult laparoscopic cholecystectomy and conversion, probably due to dense adhesions at calot’s triangle

and gall bladder fossa^{30, 31}. History of previous upper or lower abdominal surgery was not associated with difficult laparoscopic cholecystectomy in present study, which is consistent with some of the available literature^{8,23}. However, some studies reported prior history of abdominal surgery as a risk factor³².

Previous abdominal surgery may lead to adhesions between viscera or omentum and abdominal wall. There may be chances of injury to these structures during insertion of first port and risk of conversion^{14, 24}. Thickened gall bladder wall as a predictor of difficult laparoscopic cholecystectomy in this study and these results are consistent with the literature^{16, 33}. Gallbladder wall thickness greater than 3mm is suggestive of cholecystitis³³. Presence of pericholecystic collection signifies acute cholecystitis and also makes visualization of the gall bladder difficult. Presence of pericholecystic collection was not identified as a risk factor for difficult laparoscopic cholecystectomy in this study. However, in literature pericholecystic collection was identified as a significant risk factor for difficult laparoscopic cholecystectomy. Stone impaction of Hartmann's pouch was identified as a risk factor for difficult gall bladder in this study and these results are consistent with literature. Stone impaction at the neck or Hartman's pouch results hindrance in holding of the gallbladder during dissection^{23, 32}.

Based on intra-operative findings of the operating surgeon, gall bladder adhesions were noted to be associated with difficult laparoscopic cholecystectomy which is quite consistent with the literature²⁶.

Difficulty in gall bladder grasping was associated significantly with contracted gall bladder, distended gall bladder. A distended gall bladder or a gall bladder filled with stones is not easily grasped because it tends to slip away. Presence of inflammation around the gall bladder makes the wall friable and oedematous, thus posing problems to grasping.

5. CONCLUSION

The risk of difficult laparoscopic cholecystectomy increases with age. Also, prior hospitalization for acute cholecystitis is clinical risk factor associated with difficult laparoscopic cholecystectomy. On imaging studies, thick gall bladder wall and impacted stone in the Hartmann's pouch are risk factors for difficult laparoscopic cholecystectomy. The preoperative score (Randhawa) has high sensitivity, specificity, and accuracy in predicting difficult cases and can be used as a benchmark for identifying difficult laparoscopic cholecystectomies.

6. ACKNOWLEDGEMENT

The authors thank Dr. S F Kammar, Director and Dr. Gurushantappa Y, Principal KMCR for conducting this research smoothly.

7. REFERENCES

1. Bourgouin, S., Mancini, J., Monchal, T., Calvary, R., Bordes, J., & Balandraud, P. (2016). How to predict difficult laparoscopic cholecystectomy? Proposal for a simple preoperative scoring system. *American Journal of Surgery*, 212(5), 873–881. <http://doi.org/10.1016/j.amjsurg.2016.04.003>
2. Salky BA, Edye MB. The difficult cholecystectomy: problems related to concomitant diseases. *Semin Laparosc Surg*. Jun 1998;5:107-114.
3. Laws HL. The difficult cholecystectomy: problems during dissection and extraction. *Semin Laparosc Surg*. Jun 1998;5:81-91.
4. Simorov A, Ranade A, Parcels J, *et al*. Emergent cholecystostomy is superior to open cholecystectomy in extremely ill patients with acalculous cholecystitis: a large multicenter outcome study. *Am J Surg*. Dec 2013;206:935-940. Discussion 940-931.
5. Abdel Baki NA, Motawei MA, Soliman KE, Farouk AM. Pre-operative prediction of difficult laparoscopic cholecystectomy using clinical and ultrasonographic parameters. *JMRI* 2006;27(3):102-7.
6. Sahu SK, Agrawal A, Sachan PK (2013) Intraoperative Difficulties in Laparoscopic Cholecystectomy. *Jurnalul de Chirurgie (Iasi)* 2: 149-155.
7. Randhawa, J. S., & Pujahari, A. K. (2009). Preoperative prediction of difficult lap chole: A scoring method. *Indian Journal of Surgery*, 71(4), 198-201. <https://doi.org/10.1007/s12262-009-0055-y>
8. Sugrue M, Sahebally SM, Ansaloni L, Zielinski MD. Grading operative findings at laparoscopic cholecystectomy-a new scoring system. *World Journal of Emergency Surgery*. 2015 Dec;10(1):14.
9. De Mestral C, Rotstein OD, Laupacis A, Hoch JS, Zagorski B, Alali AS, *et al*. Comparative operative outcomes of early and delayed cholecystectomy for acute cholecystitis: a population-based propensity score analysis. *Ann Surg*. 2014;259:10-5.
10. Gadacz TR. Update on laparoscopic cholecystectomy, including a clinical pathway. *Surg. Clin. North Am* 2000;80:1127-45
11. Kaafarani HM, Smith TS, Neumayer L, Berger DH, Depalma RG, Itani KM. Trends, outcomes, and predictors of open and conversion to open cholecystectomy in Veterans Health Administration hospitals. *Am J Surg*. 2010;200(1):32-40.
12. Gholipour C, Fakhree MBA, Shalchi RA, Abbasi M. Prediction of conversion of laparoscopic cholecystectomy to open surgery with artificial neural networks. *BMC Surg*. 2009;9(1).
13. Tang B, Cuschieri A. Conversions during laparoscopic cholecystectomy: risk factors and

- effects on patient outcome. *J Gastrointest Surg.* 2006;10(7): 1081-1091.
14. Hussain A (2011) difficult laparoscopic cholecystectomy: current evidence and strategies of management. *Surg Laparosc Endosc Percutan Tech* 21: 211-7.
 15. Ashfaq, A., Ahmadih, K., Shah, A. A., Chapital, A. B., Harold, K. L., & Johnson, D. J. (2016). The difficult gall bladder: Outcomes following laparoscopic cholecystectomy and the need for open conversion. *American Journal of Surgery*, 212(6), 1261-1264. <https://doi.org/10.1016/j.amjsurg.2016.09.024>.
 16. Kidwai R, Pandit R, Issrani R, Prabhu N. Assessment of Risk Factors for Conversion from Difficult Laparoscopic to Open Cholecystectomy- A Hospital Based Prospective
 17. Lo CM, Fan ST, Liu CL, Lai Edward CS, Wong J. Early decision for conversion of laparoscopic to open cholecystectomy for treatment of acute cholecystitis. *Am J Surg* 1997; 173(6): 513-7.
 18. Cox MR, Wilson TG, Luck AJ, Jeans PL, Padbury RTA, Toouli J. Laparoscopic cholecystectomy for acute inflammation of gallbladder. *Ann Surg* 1993; 218(5): 630-4.
 19. Kum CK, Goh PMY, Isaac JR, Tekant Y, Ngoi SS. Laparoscopic cholecystectomy for acute cholecystitis. *Br J Surg* 1994; 81(11): 1651-4.
 20. Sippey M, Grzybowski M, Manwaring ML, *et al.* Acute cholecystitis: risk factors for conversion to an open procedure. *J Surg Res.* 2015;199(2):357-361.
 21. Yol S, Kartal A, Vatansev C, Aksoy F, Toy H. Sex as a factor in conversion from laparoscopic cholecystectomy to open surgery. *JLSL* 2006;10:359-63.
 22. Russell JC, Walsh SJ, Reed-Fourquet L, Mattie A, Lynch J. Symptomatic cholelithiasis: A Different disease in men? *Ann Surg* 1998;227:195-200.
 23. Gupta N, Ranjan G, Arora MP, Goswami B, Chaudhary P, Kapur A, Kumar R, Chand T. Validation of a scoring system to predict difficult laparoscopic cholecystectomy. *International Journal of Surgery.* 2013 Nov 1;11(9):1002- 6.
 24. Nachnani J, Supe A. Pre-operative prediction of difficult laparoscopic cholecystectomy using clinical and ultrasonographic parameters. *Indian J Gastroenterol* 2005;24:16-8.
 25. Hussien M, Appadurai IR, Delicata RJ, Carey PD. Laparoscopic cholecystectomy in the grossly obese: 4 year's experience and review of literature. *HPB (Oxford)* 2002;4:157-61.
 26. Vivek MA, Augustine AJ, Rao R. A comprehensive predictive scoring method for difficult laparoscopic cholecystectomy. *Journal of minimal access surgery.* 2014 Apr;10(2):62.
 27. Sharma, D., Kishore, K. N., Gondu, G. R., Thumma, V.M., Gunturi, S. V., Reddy, J. M., & Bheerappa, N. (2018). Predictive factors for conversion from laparoscopic to open cholecystectomy: an institutional study, 5(8), 2894-2898.
 28. Van Der Steeg HJJ, Alexander S, Houterman S, Slooter GD, Roumen RMH. Risk factors for conversion during laparoscopic cholecystectomy -experiences from a general teaching hospital. *Scand J Surg.* 2011;100(3):169-173
 29. Mirza SM, Al Salamah SM. Risk prediction for conversion from laparoscopic to open cholecystectomy. *Emir Med J.* 2003;21(2):140-145
 30. Bouarfa L, Schneider A, Feussner H, Navab N, Lemke HU, Jonker PP, *et al.* Prediction of intraoperative complexity from preoperative patient data for laparoscopic cholecystectomy. *Artif. Intell. Med* 2011 Jul;52(3):169-76
 31. Majeski J. Significance of preoperative ultrasound measurement of gallbladder wall thickness. *Am Surg.* 2007;73(9):926-9.
 32. Goonawardena J, Gunnarsson R, De Costa A. Predicting conversion from laparoscopic to open cholecystectomy presented as a probability nomogram based on preoperative patient risk factors. *Am J Surg.* 2015;210(3):492-500.
 33. Rizvi SA, Ali SA, Akhtar S, Faridi S, Ahmad M. Forecast of difficult Laparoscopic cholecystectomy by Sonography: An added advantage.
 34. Sharma, D., Kishore, K. N., Gondu, G. R., Thumma, V.M., Gunturi, S. V., Reddy, J. M., & Bheerappa, N. (2018). Predictive factors for conversion from laparoscopic to open cholecystectomy: an institutional study, 5(8), 2894-2898.
 35. Van Der Steeg HJJ, Alexander S, Houterman S, Slooter GD, Roumen RMH. Risk factors for conversion during laparoscopic cholecystectomy-experiences from a general teaching hospital. *Scand J Surg.* 2011;100(3):169-173.
 36. Mirza SM, Al Salamah SM. Risk prediction for conversion from laparoscopic to open cholecystectomy. *Emir Med J.* 2003;21(2):140-145
 37. Bouarfa L, Schneider A, Feussner H, Navab N, Lemke HU, Jonker PP, *et al.* Prediction of intraoperative complexity from preoperative patient data for laparoscopic cholecystectomy. *Artif Intell Med* 2011 Jul;52(3):169-76
 38. Majeski J. Significance of preoperative ultrasound measurement of gallbladder wall thickness. *Am Surg* 2007;73(9):926-9.
 39. Goonawardena J, Gunnarsson R, De Costa A. Predicting conversion from laparoscopic to open cholecystectomy presented as a probability nomogram based on preoperative patient risk factors. *Am J Surg.* 2015;210(3):492-500.
 40. Rizvi SA, Ali SA, Akhtar S, Faridi S, Ahmad M. Forecast of difficult Laparoscopic

DOI: 10.69605/ijlbpr_14.3.2025.135

cholecystectomy by Sonography: An added
advantage.