

ORIGINAL RESEARCH

A prospective study to evaluate the use of locking compression plate in patients of distal femur fracture managed surgically

Dr. Dhruv Gautam¹, Dr. Sunil Kumar Dhanger², Dr. Rahul Singhoria³, Dr. Bidisha Barua⁴

¹Senior Resident, ²Assistant Professor, ^{3,4}Postgraduate Student, Department of Orthopaedics, Pt. B.D. Sharma PGIMS, Rohtak, Haryana, India

Corresponding Author

Dr. Dhruv Gautam

Senior Resident, Department of Orthopaedics, Pt. B.D. Sharma PGIMS, Rohtak, Haryana, India

Email: dhruvs377@gmail.com

Received Date: 23 February, 2024 Accepted Date: 28 March, 2024

ABSTRACT

Introduction: Distal femur fracture are being treated with distal femur nail, dynamic condylar screw and even addition of a medial plate with good clinical results. By this study, we explore the potentiability of a distal femur locking plate to counter distal femur fractures of extraarticular, partial or intra-articular nature. Positive results have been brought out by various groups from all over the world. **Aim:** To study the clinical, functional and radiological outcome of distal femoral fractures managed by open reduction and internal fixation with distal femur locking plate in skeletally mature patients. **Materials and Methods:** This was a prospective study conducted from May 2023 to January 2024 in Pt. B.D. Sharma PGIMS, Rohtak, Haryana, India in skeletally mature patients with post-traumatic distal femur fractures. Follow and evaluation was done according to the Neer scoring system. **Results:** Following all principles of fracture reduction, union was achieved in all patients with mean time to radiological union being 19 weeks. The mean Range of Motion (ROM) was 109 degrees with 20 patients having a Neer score graded as excellent to satisfactory. Three patients developed complications in the form of infection (two cases) and mal-union (one case) during the course of our study, but were completely treated by the end of the study. **Conclusion:** Positive results can be obtained by distal femur locking plate alone as it is the main implant of choice for distal femur fractures of all varieties. Best outcome is expected if fracture fixation is done following all the basic principles of fracture fixation and taking benefit of the mechanical properties of a locking plate.

Keywords: Locking compression plate, Distal femur, Fracture, Surgically

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

INTRODUCTION

Distal femoral fractures account for 4-7% of all femoral fractures.¹ The distal femoral fracture occurs in approximately 37 per 1,00,000 per years.^{2,3,4} This fracture occurs in a particular bimodal distribution, 1st group patients are below 40 years of age, mostly males, sustained high energy trauma such as RTA or a fall from heights; second group consists of patients above 50 years, mostly females, with osteoporosis sustained relatively low energy trauma.^{5,6,7,8} These fractures are difficult to treat because of osteoporosis, severe comminution, too short distal femoral fragment⁹ compound injuries and associated ligament injuries, meniscal and extensor mechanism injuries.¹ The goals of surgical treatment should be anatomic reduction of the articular surface, restoration of limb alignment, length, and rotation, bone grafting for extensive bone loss and stable fixation that allows for early mobilization¹⁰ After the development of various

implants and techniques, the management of distal femoral fractures has shifted from non-operative to operative methods.^{11,12} Operative methods include traditional plating technique that require compression of the implant to the femoral shaft (blade plate, dynamic condylar screw, nonlocking condylar buttress plate), antegrade/retrograde nail fixation, submuscular locked internal fixation and external fixation.^{2,13,14,15} Distal femoral locking compression plate allows both locking and compression screw fixation of the distal femur fracture along with articular surface realignment.¹⁰ It works on the principle of rigid fixation^{2,16,17,18}, acts as an extramedullary load bearing device^{5,19} leading to early union along with knee mobilization. It works on splinting rather than compression, therefore it provides flexible stabilization causes induction of callus formation and it avoids stress.^{2,20,21} Recent generation of distal femoral locking compression plate

is precontoured plate based on the anatomy of bone. The pull out strength of locked screws better than conventional screws and that's why, it is useful in osteoporotic bone.²²

The purpose of this study is to evaluate functional outcome, fracture healing and the complications of distal femoral fractures using open reduction and internal fixation with distal femoral locking compression plate.

MATERIAL AND METHODS

This was a prospective study conducted from May 2023 to January 2024 in Pt. B.D. Sharma PGIMS, Rohtak, Haryana, India. The study was approved by Institutional Ethical Committee and informed consent was taken from all the 37 patients included in the study.

Inclusion Criteria

- 1) All patients who are above the age of 18 years having distal femoral fractures.

Exclusion Criteria

- 1) Patients of age less than 18 years.
- 2) Open distal femur fractures.
- 3) Pathological fractures.
- 4) Associated tibia fractures.
- 5) Distal femur fractures with neuro-vascular compromise.
- 6) Patient having other fractures in the same limb, hip and vertebra.

Initial resuscitation of patients was done in emergency department, limb splinted, appropriate radiographs and CT scan as per requirements was done. Preoperative investigations were done consisting of haemoglobin level, platelet count, serum electrolytes and renal function tests, coagulogram, fasting or random blood sugar levels and viral markers consisting of Hepatitis B, C and HIV. A blood group cross-match of the patient was sent to the transfusion medicine department of the hospital prior to surgery. A chest X-ray and ECG were done in those patients as required by the anaesthetic team. Spinal or epidural anaesthesia was given. Scrubbing painting draping done using standard protocols. Standard lateral approach by developing the plane between vastus lateralis muscle and lateral intermuscular septum was used for closed fractures of distal femur. Lateral parapatellar arthrotomy by using the Swash- buckler approach or its modifications were used to address the articular involvement of the lateral femoral condyle or the inter- condylar notch. A dual incision (standard lateral plus a medial subvastus incision) was also done in cases where a Hoffa's fracture or articular fracture of the medial femoral condyle had to be addressed in cases where access via the lateral approach was not

possible. We always strived for obtaining anatomical reduction and good solid fixation in the articular area of the distal femur even at the cost of spending an extra amount of time, rather than giving the patient an incongruous joint. Distal femur locking condylar plate was used for fracture fixation using locking, 4.5mm cortical screws and partially- threaded cancellous screws of diameter 4.5mm, Herbert screws for smaller articular fragments and 6.5mm for articular reduction of condyles. Technique of counter sinking for screws heads was used when proper seating of the plate had to be obtained. Length of the plate used was judged based on the extent of fracture of distal femur. In cases of fractures with metaphyseal-comminution with no bony reference point along the lateral cortex for direct fracture reduction the shaft of femur was aligned in the center of the condyles and it was ensured to prevent lateralization of the shaft at time of plate application Primary bone grafting was done in closed cases where the demand for such a procedure was gauged by the operating surgeon. The average blood loss was 300- 400ml in each case. No use of tourniquets was done in any of the case. Post-operatively suction drain was removed after 24-48 hours and first wound inspection was done on 2nd post-operative day. Intra-venous antibiotics were continued for 24 hours in closed fractures. Post-operative physiotherapy regime was tailored according to the fracture pattern and fixation achieved. In cases where ever possible, knee bending and assisted knee Range of Motion (ROM) exercises were initiated on postoperative day 1. Patients were mobilized based on the degree of bone quality, severity of injuries, and pattern of fractures. At post-operative day 2 to 3, the patients were mobilized with crutches/walker until 6 weeks. Full weight bearing ambulation without any aids was started at approximately 3 months in majority of the cases with radiographic evidence of fracture union. Patients were discharged at post-operative day 3 with stitch removal being done on day 14th, making it convenient for the patient to take bath and maintain good body hygiene. Regular follow-up of all cases was done at 6 weeks, 12 weeks and 24 weeks. At each follow-up patients were evaluated clinically using knee society Neer's scoring system and radiologically with appropriate x-rays. No patients were lost to follow-up.

Statistical analysis

The statistical data analysis was carried out using a computer based statistical analysis programme, SPSS version 20 (IBM, Chicago, USA). For the statistical data analysis, paired t-test was used between two correlated groups while for uncorrelated groups; means were compared using independent t-test. A p-value of <0.05 was considered as significant.

RESULTS

Table 1: Distribution of age of distal femur fracture patients

Age (year)	Frequency (N)	Percent (%)
20-30	3	8.1
30-40	12	32.4
40-50	8	21.6
50-60	5	13.5
≥60	9	24.3
Total	37	100

Out of the 37 cases in our study, the mean age of cases in our study was 45.70±15.35 years ranging from 20 to 78 years approximately 25(67.6%) of the patients were male, 12(32.4%) were female. The road traffic accident was the most common mode of injury,

in 27 number (72.9%). Fall due to various reasons experienced by 10 patients (27.1%) The side affected in distal femur fracture among studied patients was noted more on left by 54.1% while the right seventeen (45.9%) cases.

Table 2: The frequency and percentage distribution of preoperative diagnosis to muller’s classification

Preoperative Diagnosis	Frequency (N)	Percent (%)
Muller Type A-1	10	27.0
Muller Type A-2	3	8.1
Muller Type A-3	5	13.5
Muller Type B-1	7	18.9
Muller Type B-2	2	5.4
Muller Type B-3	2	5.4
Muller Type C-1	4	10.8
Muller Type C-2	1	2.7
Muller Type C-3	3	8.1

In our study, 6(16.2%) patient had post operative complication two patients had early post-operative complication in the form of superficial infection. These were then treated with culture sensitive parenteral antibiotics and antiseptic dressing, which resulted in satisfactory healing of the fracture. One case was found to have late complication in the form of mal-union of the distal femur fracture [Table 3].

Out of total, 37 patients, six patients (16%) achieved full flexion of 130 degrees or more. Better outcome was observed in terms of range of motion at knee joint in extra-articular fractures (82%) than intra-articular fractures (72%) treated with distal femur locking plate. The mean ROM in all 37 patients was 114 degrees.



Figure 1 clinical photograph at 6 month followup



Figure 2 clinical photograph at 6 month followup

Table 3: Distribution of late complications of distal femur fracture patients


Late complications	Frequency (N)	Percent (%)
Wound infection	2	5.4
Knee stiffness	3	8.1
Malunion	1	2.7
Total	6	16.2

According to Neer’s criteria which included evaluation in terms of pain, range of motion, walking


and work capacity, anatomy and X-ray findings score at 6 months was calculated results were compared in

case of intra- and extra-articular fractures, clinically ROM at knee joint and score at 6 months was better in extra-articular fractures than intra-articular, although statistically insignificant (p-value = 0.5012) in this situation as well. In the present study, results were

tabulated into 4 groups i.e., excellent, satisfactory, unsatisfactory and poor according to criteria laid down by Neer et al., [11], the results were excellent in 14 (37.8%), good in 8 (48.6%) fair in 3 (8.1%) and poor in 2 (5.4%) patients.



Pre-Operative X-ray Antero-Posterior view and lateral view



Post operative X ray Anteroposterior AND Lateral view

Table 4: Assessment of clinical and functional outcome by neer’s society scoring system

Scoring with Category		AT First Follow up		At second Follow up		At Third Follow up	
		n	%	n	%	n	%
<55	Poor	32	86.5	10	27.0	2	5.4
55-69	Fair	4	10.8	12	32.4	3	8.1
70-85	Good	1	2.7	15	40.5	18	48.6
>85	Excellent	0	0.0	0	0.0	14	37.8
Total		37	100.0	37	100.0	37	100.0
Mean±Std. Deviation		43.54±10.65 points		63.11±12.50 points		79.03±12.15 points	




Table 5: Distribution of time of union of distal femur fracture patients

Time of union	Frequency (N)	Percent (%)
11-13 weeks	7	18.9
13-15 weeks	11	29.7
15-17 weeks	11	29.7
17-19 weeks	8	21.6
Total	37	100

Most of the cases (44%) attained radiological union in 19-22 weeks. The mean time for radiological union was 19 weeks

DISCUSSION

The mean age in our study was 45.70 years . A study published by Reddy J.A.V mean age was 40 years (range 20-70) ²³ Ramanand M. et al found mean age 52.25 years (range 20-60 years)²⁴

Dr. Pugazhendhi et al conducted a short term prospective study of the functional outcome in distal femoral fractures treated by locking compression plate, they reported that out of 22 patients, there were 12 patients in 21-30 age group, 5 patients in 31-40 age group, 3 patients in 41-50 age group and 2 patients in 51-60 age group. They concluded that the most common age group was 21-30 years.²⁵. Scope of distal

femur locking plate is limited not only to isolated distal femur fractures but also in the use of peri-prosthetic distal femur fractures in patients of Total Hip Replacement (THR) ²⁶and Total Knee Replacement (TKR). Even extreme distal peri-prosthetic supracondylar fractures can be managed with lateral locked plate with predictable results similar to those seen in more proximal fractures²⁷

In our study, the average time of union was 14.5 weeks, with majority of cases united between 13-17 weeks. Time of union in other studies are tabulated below in table

Author	Time of union (weeks)
Shreedhar M. et al ²⁸	20
Pugazhendhi G. et al ²⁵	16
Reddy J.A.V. et al ²³	12
Shriharsha et al ²⁹	19.6
Panchal P. et al ³⁰	18

Ramesh Krishna et al ¹	15.3
Mahesh D.V. et al ¹⁰	18
Our study	14.3

In our study, by using Neer's society scoring system, the results were excellent in 14 (37.8%), good in 18 (48.6%), fair in 3 (8.1%) and poor in 2 (5.4%) patients. Our study was comparable with standard

studies of management of distal femur fracture using distal femoral locking compression plate. Functional outcome of other studies are tabulated below in table

Author	Results			
	Excellent	Good	Fair	Poor
Ramesh Krishna et al ¹	15	11	3	1
Tapi Nalo et al ³¹	25	3	1	0
C.V. Dasaraiah et al ²	15	11	3	1
Pradip B Patil et al ³²	18	11	1	0
Shriharsha R.V. et al ²⁹	4	10	5	6
Parth Panchal et al ³⁰	11	5	3	1
Our study	14	18	3	2

Dr. Pugazhendhi G. et al concluded that, the early complications were encountered in 3 patients, 2 patients developed superficial infection and 1 patient had wound gaping. Late complications included knee stiffness in 2 patients. None of the patients had loss of fixation or malunion or non union.²⁵ According to Parth Panchal³⁰ et al study on treatment of distal end of fracture femur by locking compression plate, 5 patients developed complications, with 4 (20%) patients experiencing superficial infection and 1 (5%) patients delayed union. Tapi Nalo et al conducted the study on treatment of supracondylar fracture of distal femur with condylar locking compression plating, and they reported that, there were 2 cases of superficial infection.³¹

LIMITATIONS

The requirement for more number of patients was felt during the conduct of our study but with a longer follow-up of 2 years it was reasonable to restrict ourselves to a follow-up compliant group.

CONCLUSION

Distal femur locking plate is still the way forward for treating distal femur fractures. Positive results have been published by researchers with implants such as distal femur nail, dynamic condylar screw and even addition of a medial plate to a distal femur locking plate for treating distal femur fractures. By conducting this study we can put forward the assumption with some degree of confidence that fractures of the distal femur of all varieties extra-articular, partial articular and intra-articular non-comminuted as well as comminuted ones, if fixed in a proper fashion following all the basic principles of fracture fixation, good results can be obtained by using a distal femur locking plate alone as the main implant of choice for such fractures. It has also reduced the requirement of secondary procedures such as bone grafting especially in fractures with metaphyseal comminution

REFERENCES

1. Krishna RK, Nayak MA. Study of surgical management of distal femoral fracture by distal femoral locking compression plateosteosynthesis. *IJOS* 2015;1(1):22-6
2. Dasaraiah CV, Rao AS. Study of surgical management of supracondylar femoral fracture by locking compression plate. *IOSR J Dent Med Sci (IOSR-JDMS)* 2016;15 (2):23-33.
3. Martinet O, Cordey J, Harder Y, Maier A, Buhler M, Barraud GE. Epidemiology of fracture of distal femur. *Injury* 2000;31:62-63.
4. Arneson TJ, Melton LJ III, Lewallen DG, O'Fallon WM. Epidemiology of diaphyseal and distal femoral fractures in Rochester, Minnesota, 1965-1984. *Clin Orthop* 1988;234:188-94
5. Gupta SKV, Govindappa CVS, Yalamanchili RK. Outcome of retrograde intramedullary nailing and locking compression plating of distal femoral fractures in adults. *Open access Orthopaedics* 2013;1(3):23.
6. Verma R, Kumar K, Sidhu V, Saraf A. An overview of cases of fracture distal femur: a hospital based study. *Int J Med Res Prof* 2018;4(6):319-22. DOI:10.21276/ijmrp.2018.4.6.074
7. Crist B, Rocca G, Murtha Y. Treatment of acute distal femur fractures. *Orthopaedics* 2008; 31:681.
8. Higgins TF. Distal femoral fractures. *J Knee Surg* 2007;20:56-66.
9. Hakeem A, Khan N, Khan M, Ullah F. Dynamic condylar screw fixation in treatment of supracondylar fracture of distal femur. *Rawal Med J* 2010;35(1):45-47.
10. Mahesh DV, Gunnaiah, Vishwanath. Management of distal femur fracture by locking compression plate. *IJHSR* 2014;4(5):235-40
11. Yeap EJ, Deepak AS. Distal femoral locking compression plate fixation in distal femoral fractures: early results. *Malaysian Orthop J* 2007;1(1):12-7.
12. Schatzker J, Lambert DC. Supracondylar fractures of the femur. *Clin Orthop* 1979;138:77-83.
13. Wilson JN. Watson Jone's: Fractures and joint injuries. 6th ed, 1982. p. 1003-070, 1982.
14. Neer CS II, Grantham SA, Shelton ML. Supracondylar fracture of the adult femur - a study of one hundred and ten cases. *JBJS Am* 1967;49-A(4): 591-613.

15. Brown A, D'Arcy JC. Internal fixation for supracondylar fractures of the femur in the elderly patient. *JBSJ* 1971;53B(3):420-424
16. Thomas PR, Richard EB, Christopher GM. *A O Principles of Fracture Management*. 2nd edition. Switzerland: AO Publishing; 2007.
17. Krettek C, Schandelmaier P, Miclau T, Tscherner H. Minimally invasive percutaneous plate osteosynthesis (MIPPO) using the DCS in proximal and distal femoral fractures. *Injury* 1997; 28(1): 20- 30.
18. Kregor PJ, Stannard J, Zlowodzki M, Cole PA, Alonso J. Distal femoral fracture fixation utilizing the Less Invasive Stabilization System (L.I.S.S.): The technique and early results. *Injury* 2001; 32(3): 32-47
19. Smith WR, Ziran BH, Anglen JO, Stahel PF. Locking plates: tips and tricks. *J Bone Joint Surgery Am* 2007;89(10):2298-307.
20. Venable CS, Stuck WG, Beach A. The effects on bone of the presence of metals: based on electrolysis. *Ann Surg* 1937 June.
21. Kregor PJ, Stannard JA, Zlowodzki M, Cole PA. Treatment of distal femur fractures using the less invasive stabilization system: surgical experience and early clinical results in 103 fractures. *J Orthop Trauma* 2004;18(8):509-20
22. Kumar GNK, Sharma G, Farooque K, Sharma V, Ratan R, Yadav S, Lakhota D. Locking compression plate in distal femoral intraarticular fractures: our experience. *Int Scholarly Res Notices* 2014; Article ID 372916, 5 pages.
23. Reddy JAV, Chary NB. Functional outcome of distal end of femur with locking compression plate: a prospective study. *Ann Int Med Den Res* 2016;2(1):362-65.
24. Ramanand M, Mudgal CV, Nagendrappa M. Management of distal femoral fractures- A comparative study between supracondylar nail and distal femoral locking plate. *J Evid Based Med Healthc* 2017;4(34),2073-2077.
25. Pugazhendhi G, Venkatachalam K, Madhukar. A short term prospective study of the functional outcome in distal femoral fractures treated by locking compression condylar plates. *Sch Acad J Biosci* 2016; 4(9):707-11
26. Chakravarthy J, Bansal R, Cooper J. Locking plate osteosynthesis for Vancouver Type B1 and Type C periprosthetic fractures of femur: a report on 12 patients. *Injury* 2007;38(6):725-33.
27. Streubel PN, Gardner MJ, Morshed S, Collinge CA, Gallagher B, Ricci WM, et al. Are extreme distal periprosthetic supracondylar fractures of the femur too distal to fix using a lateral locked plate? *J Bone Joint Surg Br* 2010;92(4):527-34.
28. Sreedhar M, Luther ME, Anudeep B, Prashanth, Swaroopa, Madhav V. Functional and radiological outcome of distal femoral fractures treated with locking compression plate. *J Evol Med Dent Sci* 2015;4(38):6664-71.
29. Shriharsha RV, Sapna M. Utility and outcomes of locking compression plates in distal femoral fractures. *Int J Res Orthop* 2015;1(1):15-21.
30. Panchal P, Patel C, Poptani A. Treatment of distal end of fracture femur by locking compression plate. *Int J Med Sci Pub Hlth* 2016;5:09.
31. Nalo T, Agrahari A, Datta S, Paul V, Singh SN, Langshong R. Treatment of supracondylar fracture of distal femur with condylar locking compression plating. *Int J Sci Res (IJSR) ISSN (Online): 2319-7064*.
32. Patil PB, Patil RS, Lad SA, Suren T, Jain UD. The study of functional outcome of distal end femur fractures operated with locking compression plate. *Asian Pac J Health Sci* 2016;3(1):135-9.