

**ORIGINAL RESEARCH**

# Acinetobacter Meningitis: A Retrospective Study on its outcome and Antimicrobial resistance pattern in Post neurosurgical meningitis Patients at a Tertiary Care Centre

<sup>1</sup>Dr. Anjum Ara Mir, <sup>2</sup>Dr. Sayim Wani, <sup>3</sup>Dr. Nargis Bali, <sup>4</sup>Dr. Munaza Aman, <sup>5</sup>Dr. Susheem R

<sup>1,2,4</sup>Senior Resident, <sup>3</sup>Associate Professor, <sup>5</sup>Post graduate student, Department of Microbiology, Sher-i-Kashmir Institute of Medical Sciences, Soura, Srinagar, Jammu and Kashmir, India

**Corresponding author**

Dr. Munaza Aman

Senior Resident, Department of Microbiology, Sher-i-Kashmir Institute of Medical Sciences, Soura, Srinagar, Jammu and Kashmir, India

Email: [drmunaza27aman@gmail.com](mailto:drmunaza27aman@gmail.com)

Received: 18 April, 2024

Accepted: 6 May, 2024

**ABSTRACT**

**Introduction:** *Acinetobacter baumannii* (*A. baumannii*) is a gram negative bacillus and mortality rate due to *Acinetobacter baumannii* nosocomial meningitis (ANM) is high. The increasing antimicrobial resistance makes the treatment options available very limited necessitating the need for a continuous Antimicrobial Resistance surveillance that would aid in the proper implementation of the empirical therapy while taking into consideration its extensive antibiotic resistance spectrum. **Objectives:** This study aims to determine the prevalence & antibiotic resistance pattern of *A. baumannii* isolated from the CSF samples from the post neurosurgical cases. **Material and methods:** A cross-sectional study was conducted over two years, from December 2021 to November 2023, at the Department of Microbiology, Sher-i-Kashmir Institute of Medical Sciences, Srinagar. Data from 28 patients with positive CSF samples for *A. baumannii* were collected, including age, gender, antimicrobial susceptibility, and mortality. CSF samples were received in culture bottles and incubated in the BacT/ALERT system. Positive bottles underwent gram staining and subcultures on blood, MacConkey agar, and Chocolate agar. Non-lactose fermenter colonies were identified using VITEK Densichek, and antimicrobial susceptibility testing was conducted using the VITEK 2 ID-GNB card and AST-NO09 card respectively in VITEK 2 system with software release 2.01. **Results:** Mean age of the patients with *Acinetobacter meningitis* included in this study was 32.2 years. 16/28 (57.1%) were male. All cases were associated with previous neurosurgical procedures. Of the total number of patients enrolled in the study, 11 patients had MDR isolates whereas 2 patients had XDR *Acinetobacter*. 19 (67.8%) and 25 (89.2%) of the isolates were resistant to Imipenem and Meropenem respectively. Mortality rate of the patients was 60.7%. **Conclusion:** In critically ill patients post-neurosurgery, *A. baumannii* infection poses a significant challenge. Understanding the pattern and antibiotic susceptibility of *Acinetobacter meningitis* is crucial for effective antimicrobial strategies, reducing mortality and morbidity. Continued efforts are required to develop improved antimicrobial policies and implement effective infection control practices. Given the high mortality rates observed, aggressive empirical treatment, including intrathecal therapy, is necessary in regions with endemic meningitis.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**INTRODUCTION**

*Acinetobacter baumannii* (*A. Baumannii*) is a pleomorphic, non-motile gram negative bacillus that belongs to the genus *Acinetobacter*, which in turn has as many as 30 species. Most of these species are found in the environment and are not pathogenic to human beings. The most commonly reported species are termed as *A. calcoaceticus*-*A. Baumannii* (ACB)

complex which includes *Acinetobacter calcoaceticus* and *A. baumannii*. Among these, *A. baumannii* is clinically the most significant. It is an opportunistic pathogen owing to its ability to cause higher rate of infection in immunocompromised individuals. In the recent past, it has been termed as a "red alert" human pathogen.[1,2] It is known to cause a large number of infections including infections involving the

respiratory tract, bacteremia, meningitis, and wound infection. [3] Patients undergoing neurosurgical procedures are very much prone to develop Hospital-acquired meningitis caused by a wide range of Gram negative and Gram positive microorganisms among which the majority of nosocomial meningitis is caused by *A. baumannii*. [4] Craniocerebral operations put the patients at a higher risk of suffering from bacterial meningitis caused by *A. baumannii* with extremely fatal consequences. *A. baumannii* meningitis has been seen to be associated with external ventricular drainage (EVD), cerebrospinal fluid (CSF) leaking, or head trauma, Ventriculo-Peritoneal (VP) shunt and is the most common pathogen isolated in this subset of patients [4,5]. About 4% of all meningitis and shunt-related infections are caused by *A. baumannii* and these are known to show mortality rates nearing to 70%. The transmission of this pathogen occurs through the vicinity of affected patients or colonizers such as linens fomites, curtains, bed rails, tables, doors, feeding tubes, and even medical equipment and it is able to survive desiccation and remain viable for months which again is an important factor that facilitates its spread in the hospital. In addition to this, Contaminated respiratory support equipment, suction devices, and devices used for intravascular access also serve as an important source of infection. [6] Also, age over 40, raised cerebrospinal fluid white blood cell count, and diabetic and hypertensive patients are at a higher risk for mortality due to *A. baumannii* in the neurosurgical ICU. [7] The *A. baumannii* infections are usually treated with antibiotics like Carbapenems, Aminoglycosides, Polymyxins, Tigecycline, Tetracyclines and Combination Therapy. [8] At the same time, there has been a significant increase in the Multidrug-resistance of these isolates which is being recognized as a growing concern since resistance to all commercially available antimicrobials (aminoglycosides, cephalosporins, quinolones and imipenem) raises an important therapeutic issue. [9] In addition to all this, the penetration of antibiotic agents through the blood-brain barrier into the cerebrospinal fluid (CSF) is variable and further limits the therapeutic choices for these infections. [10]

The increasing antimicrobial resistance makes the treatment options available very limited necessitating the need for a continuous AMR surveillance that would in turn aid in the proper implementation of the empirical therapy while taking into consideration its extensive antibiotic resistance spectrum. This study aims to determine the prevalence & antibiotic resistance pattern of *A. baumannii* isolated from the CSF samples from the post neurosurgical cases.

## MATERIAL AND METHODS

### Study period

This cross-sectional study was conducted for a period of 2 years from December 2021 to November 2023 in

Department of Microbiology at the Sher-i-Kashmir institute of medical sciences, Srinagar.

## METHODOLOGY

Data of 28 patients whose CSF samples were positive for *A. Baumannii* were obtained with respect to age, gender, investigations done, and antimicrobial susceptibility pattern and mortality. Post-neurosurgical Meningitis was defined as meningitis developing within 3 months after neurosurgery and was diagnosed based on specific criteria: (1) Isolation of *A. baumannii* from cerebrospinal fluid (CSF); (2) Presence of at least one symptom or sign without alternative explanations, including fever ( $\geq 38^{\circ}\text{C}$ ), headache, vomiting, confusion, irritability, or meningeal irritation; (3) History of neurosurgical intervention within the preceding 3 months. [11,12] A patient was considered cured upon disappearance of symptoms and signs along with sterile CSF. [13]

CSF samples were received in the CSF culture bottles which were incubated at  $35 \pm 2^{\circ}\text{C}$  in the BacT/ALERT blood culture System and the bottles that showed a positive signal were subjected to microscopic examination of gram-stained smears of their contents. At the same time, Subcultures were performed from these bottles on blood, MacConkey agar plates and Chocolate agar which were incubated aerobically at  $37^{\circ}\text{C}$  overnight. A suspension was prepared from the plates that showed growth as Non lactose fermenter on Macconkey agar and the turbidity of the bacterial suspension was adjusted with VITEK Densichek (bioMérieux) to match the McFarland 0.5 standard in 0.45% sodium chloride. Afterward, the VITEK 2 ID-GNB card, the AST-NO09 card, and the bacterial suspension were manually loaded into the VITEK 2 system. The VITEK 2 system reported the results automatically with software release 2.01. Standard strains including *Escherichia coli* ATCC 25922 and *Pseudomonas aeruginosa* ATCC 27853 were used as controls.

## RESULTS

Mean age of the patients with *Acinetobacter* meningitis included in this study was 32.2 years. 16 were male. Multi drug resistant *Acinetobacter* spp. was defined as the isolates resistant to at least three classes of antimicrobial agents. Extensively Drug Resistant *Acinetobacter* spp. was defined as the isolate that is resistant to the three classes of antimicrobials and also resistant to carbapenems. Finally, 'Pan drug resistant *Acinetobacter* spp. was defined as the XDR *Acinetobacter* spp. that is resistant to polymyxins and tigecycline. [14] Of the total number of patients enrolled, 11 patients had MDR isolates whereas 2 patients had XDR *Acinetobacter* was isolated from 2 patients. Mortality rate was found to be 60.7%. There was no significant association observed between gender, length of hospital stay, and the mortality rate of the patients.

**Table 1: Gender and outcome of 28 patients with Acinetobacter meningitis**

No	Gender	Outcome
1.	Female	Survival
2.	Female	Death
3.	Male	Death
4.	Female	Survival
5.	Male	Death
6.	Female	Survival
7.	Male	Death
8.	Male	Survival
9.	Female	Death
10.	Male	Death
11.	Male	Survival
12.	Female	Death
13.	Male	Survival
14.	Male	Survival
15.	Male	Death
16.	Male	Survival
17.	Female	Death
18.	Male	Death
19.	Male	Death
20.	Female	Survival
21.	Male	Death
22.	Female	Survival
23.	Male	Death
24.	Female	Survival
25.	Female	Death
26.	Male	Death
27.	Female	Death
28.	Male	Death

**Table 2: Resistance profile of Acinetobacter isolates**

Antibiotic	N(%)
Amikacin	10(83.9%)
Gentamicin	26 (92.8)
Ampicillin	4(14.2)
Cefepime	26 (92)
Ampicillin+Sulbactam	1(3.57)
Ceftriaxone	23(82.1)
Imipenem	25(89.2)
Meropenem	19(67.8)
Piperacillin+Tazobactam	19(67.8)
Tigecycline	0
Colistin	0

**Table 3: Distribution of isolates on the basis of gender and length of stay in the hospital.**

Variable	Alive N(%)	Dead N(%)	P-value
Gender			
Male	9 (75)	7 (43.5)	0.184
Female	3 (25)	9(56.5)	
Length of hospitalization (days) median			
28 days	8(66.6)	6(37.5)	0.196
17 days	4(33.4)	10(62.5)	

## STATISTICAL ANALYSIS

All the statistical analysis was done using SPSS software.V.23. All categorical variables were shown in the form of frequency and percentage. All variables were discussed at 5% level of significance i.e. value of  $P < 0.05$ .

## DISCUSSION

Meningitis is diagnosed on the basis of clinical symptoms and/or positive cerebrospinal fluid (CSF) cultures. Various gram positive and gram negative bacteria are responsible for causing meningitis in patients who have underwent neurosurgical intervention and this life-threatening condition is associated with a large number of severe complications necessitating the need for prompt and appropriate medical treatment and *A. baumannii* is considered to be a serious public health problem owing to their prevalence and mortality in developing countries.[15,16] Among the gram negatives, *A. baumannii* is known to be the most common pathogen causing post neurosurgical procedures.[4] After craniocerebral operations in neurosurgery the patients are at a high risk to suffer from bacterial meningitis caused by *A. baumannii*. *A. baumannii* meningitis is well recognized and has been described by many doctors worldwide.[13] *Acinetobacter* spp. is able to survive in a hospital milieu and can persist for a large period of time on surfaces.[17] A great challenge for physicians and clinical microbiologists is the management of multidrug-resistant *Acinetobacter* spp. infections. Carbapenems such as meropenem or imipenem are considered first-line, however carbapenem-resistant species have emerged recently.[18]

In this study, there was no major difference between the number of patients from each gender with 42.9% patients being female and 57.1% being male. Similar findings were discovered in a study conducted by Metan et al. [19] Length of stay was 17-28 days. Comparable results were uncovered in research conducted by Metan and colleagues.[19]

Majority of the isolates were resistant to imipenem (89.2%), meropenem (85%), gentamicin (92.8%), amikacin (83.9%), ampicillin (14.2%), Ampicillin plus sulbactam (3.57%), ceftriaxone (82.1%), Piperacillin-Tazobactam (67.8%) and none of the isolates were found to be resistant to tigecycline or colistin. Similar pattern of resistance was seen in a study conducted by Kar et al. (4) Contrary to this, a study conducted by Tuon FF et al revealed a low resistance. [20]. The mortality rate in the present study was 67.8%. Similar Mortality rate of by Tuon FF et al., [20] who reported a mortality rate of 72.7% and by Sharma R et al., [6] who reported about 40% mortality. Contrary to these results, a mortality rate of 28.57%, 30% and 20%, respectively was seen in a study conducted by Sipahi OR et al.[21] The high mortality rate in our setting could be attributed to lack of strict infection control measures at our institution.

Considering the high prevalence and mortality rates of *Acinetobacter* meningitis, steps need to be taken to reduce the post-surgery or shunt infection which include following strict infection control measures.

## CONCLUSION

In critically ill people who have underwent neurosurgical intervention, a formidable challenge to proper management is infection with *Acinetobacter baumannii*. Information on the pattern and antibiotic susceptibility of *Acinetobacter* meningitis is important in modulating antimicrobial policy that would in turn help in reducing the mortality and morbidity related to infections. Hence, continued efforts are needed to develop better antimicrobial policies against this pathogen that will in turn help in implementation of appropriate infection control practices.

## REFERENCES

- Howard A, O'Donoghue M, Feeney A, Sleator RD. *Acinetobacter baumannii*: an emerging opportunistic pathogen. *Virulence*. 2012 May 1;3(3):243-50.
- Xiao J, Zhang C, Ye S. *Acinetobacter baumannii* meningitis in children: a case series and literature review. *Infection*. 2019;47:643-649.
- Raut S, Rijal KR, Khatiwada S, Karna S, Khanal R, Adhikari J, Adhikari B. Trend and characteristics of *Acinetobacter baumannii* infections in patients attending Universal College of Medical Sciences, Bhairahawa, Western Nepal: a longitudinal study of 2018. *Infect Drug Resist*. 2020 Jun 8;13:1631-1641.
- Kar M, Dubey A, Singh R, Sahu C, Singh S. *Acinetobacter* meningitis: a retrospective study on its incidence and mortality rates in postoperative patients at a tertiary care centre in Northern India. *J Clin Diagn Res*. 2023 Jan;17(1):DC01-DC06.
- Ni S, Li S, Yang N, Zhang S, Hu D, Li Q, Lu M. Post-neurosurgical meningitis caused by *Acinetobacter baumannii*: case series and review of the literature. *Int J Clin Exp Med*. 2015 Nov 15;8(11):21833-8.
- Asif M, Alvi IA, Rehman SU. Insight into *Acinetobacter baumannii*: pathogenesis, global resistance, mechanisms of resistance, treatment options, and alternative modalities. *Infect Drug Resist*. 2018 Aug 21;11:1249-1260.
- Ayoub Moubareck C, Hammoudi Halat D. Insights into *Acinetobacter baumannii*: a review of microbiological, virulence, and resistance traits in a threatening nosocomial pathogen. *Antibiotics*. 2020;9(3):119.
- Fishbain J, Peleg AY. Treatment of *Acinetobacter* infections. *Clin Infect Dis*. 2010 Jul 1;51(1):79-84.
- Kendirli T, Aydin HI, Hacıhamdioglu D, Akin R, Lenk MK, Gokcay E, et al. Meningitis with multidrug-resistant *Acinetobacter baumannii* treated with ampicillin/sulbactam. *J Hosp Infect*. 2004;56(4):328.
- Jindal N, Jain S, Bhowmick A, Bhargava V. A lurking threat of community-acquired *Acinetobacter* meningitis—a rare case report from Punjab, India. *Medicines*. 2022;9(4):27.
- Chang CJ, Ye JJ, Yang CC, Huang PY, Chiang PC, Lee MH. Influence of third-generation cephalosporin resistance on adult in-hospital mortality from post-neurosurgical bacterial meningitis. *J Microbiol Immunol Infect*. 2010;43:301-309.

12. Khan FY, Abukhattab M, Baager K. Nosocomial postneurosurgical *Acinetobacter baumannii* meningitis: a retrospective study of six cases admitted to Hamad General Hospital, Qatar. *J Hosp Infect.* 2012;80:176–179.
13. Ni S, Li S, Yang N, Zhang S, Hu D, Li Q, Lu M. Post-neurosurgical meningitis caused by *Acinetobacter baumannii*: case series and review of the literature. *Int J Clin Exp Med.* 2015 Nov 15;8(11):21833-8. PMID: 26885152; PMCID: PMC4723998.
14. Manchanda V, Sanchaita S, Singh N. Multidrug-resistant *Acinetobacter*. *J Glob Infect Dis.* 2010;2(3):291-304.
15. Schuertz KF, Tuon FF, Palmeiro JK, Conte D, Telles JPM, Trevisoli LE, Dalla-Costa LM. Bacteremia and meningitis caused by OXA-23-producing *Acinetobacter baumannii* – molecular characterization and susceptibility testing for alternative antibiotics. *Braz J Microbiol.* 2018;49(SI):199-204. DOI: 10.1016/j.bjm.2018.04.002.
16. Chojak R, Kozba-Gostyla M, Gaik M, Madei M, Majerska A, Soczynski O, Czapiga B, et al. Meningitis after elective intracranial surgery: a systematic review and meta-analysis of prevalence. *Eur J Med Res.* 2023;28:184.
17. Manchanda V, Sanchaita S, Singh N. Multidrug-resistant *Acinetobacter*. *J Glob Infect Dis.* 2010 Sep;2(3):291-304.
18. Mak J, Mittal A. Ventriculostomy-related *Acinetobacter* meningitis. *Am J Respir Crit Care Med.* 2022;205:A1617.
19. Metan G, Alp E, Aygen B, Sumerkan B. *Acinetobacter baumannii* meningitis in post-neurosurgical patients: clinical outcome and impact of carbapenem resistance. *J Antimicrob Chemother.* 2007 Jul;60(1):197–199.
20. Tuon FF, Pentead-Filho SR, Amarante D, Andrade MA, Borba LA. Mortality rate in patients with nosocomial *Acinetobacter* meningitis from a Brazilian hospital. *Braz J Infect Dis.* 2010;14(5):237-40.
21. Sipahi OR, Mermer S, Demirdal T, Ulu AC, Fillâtre P, et al. Tigecycline in the treatment of multidrug-resistant *Acinetobacter baumannii* meningitis: results of the Ege study. *Clin NeurolNeurosurg.* 2018;172:31-38.